RESTRICTED USE PESTICIDE

DUE TO HAZARDS TO NON-TARGET SPECIES

For retail sale to, and use by, Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Ramik[®] Brown

Weather-Resistant Rodenticide

For use in and around structures and in terrestrial nonfood/nonfeed areas to control commensal rats and mice and in Fruit Tree and Nut (Pecans, Almonds, Walnuts, Filberts/Hazelnuts) orchards, Christmas tree farms, tree plantations and commercial nurseries to control voles.

ACTIVE INGREDIENT:

Diphacinone

 (2-Diphenylacetyl-1,3-Indandione)
 0.005%

 INERT INGREDIENTS:
 99.995%

 TOTAL:
 100.000%

EPA Reg. No. 61282-45 EPA Est. No. 61282-WI-1

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID Have label with you when obtaining treatment advice If Call a poison control center, doctor, or 1swallowed 800-498-5743 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. If on skin or Take off contaminated clothing. clothing Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center, doctor, or 1-800-498-5743 immediately for treatment advice.

NOTE TO PHYSICIAN: If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K_1 , intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Wear gloves if bait may be handled. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to mammals and birds. Do not contaminate water when disposing of equipment washwater or rinsate. Do not apply this product directly to water, or to areas

where surface water is present or to intertidal areas below the mean high water mark.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

STRUCTURES AND TERRESTRIAL, NONFOOD/ NONFEED AREAS FOR NORWAY RAT, ROOF RAT AND HOUSE MOUSE CONTROL

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help to prevent accidents:

- Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger stations are needed in areas open to hoofed livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, and unused, spoiled and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway rats, roof rats and house mice.

IN AND AROUND STRUCTURES:

Urban areas: May only be used in and around (within 15 ft.) of homes, industrial, agricultural, public, port, and terminal buildings, and similar man-made structures. May also be used in transport vehicles (ships, trains, aircraft) and alleys.

Non-urban areas: May only be used in and around (within 15 ft.) of homes and agricultural buildings.

TERRESTRIAL, NONFOOD/NONFEED AREAS:

This product may also be used in the following areas: river banks, irrigation ditches, gullies, railroad tracks, fences, and

sanitary landfills. All placements must be in tamper-resistant bait stations or deeply into rat burrows.

Note: This weather-resistant bait may be placed at the above sites, even if wet or damp. Do not place bait in areas where there is a possibility of contaminating food/feed or surfaces that come in direct contact with food/feed. Do not broadcast bait.

SELECTION OF TREATMENT AREAS:

Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS: Wear waterproof gloves. **Norway and Roof Rats**

Place 4 to 16 oz. of bait per placement. Place bait at 15- to 30-foot intervals. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

House Mice:

Apply ¼ to ½ oz. (1-2 level tablespoons) of bait at 8- to 12-foot intervals. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

Follow-up

Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and dispose of all dead rodents and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. Where a continuous source of infestation is present, maintain permanent bait stations and replenish with census or toxic bait as needed.

FRUIT TREE ORCHARDS (BEARING AND NON-BEARING) FOR PINE AND MEADOW VOLE CONTROL

USE RESTRICTIONS: For control of pine and meadow voles in pome fruit (apples, pears) and stone fruit (peaches, nectarines) tree orchards in these states. Aerial application is only permitted in those following states where aerial is listed next to the state name.

Connecticut Michigan/aerial Georgia Missouri Idaho/aerial Montana/aerial Massachusetts North Carolina New Hampshire South Carolina Ohio/aerial Utah/aerial Oregon/aerial Virginia/aerial Pennsylvania/aerial Vermont/aerial Washington/aerial West Virginia/aerial **Timing**: Apply after fall harvest (including drops), before new spring growth, and during which three days of rain-free and snow-free weather are expected.

Grazing: Do not graze animals in treated areas.

Hay: Do not use hay cut after application for feed or bedding.

Bare Ground and Unspecified Sites: Do not apply over bare ground or crops unspecified above.

Application Type: Apply only by hand spot baiting and ground broadcasting and aerial broadcast in those states specified above.

PRE-APPLICATION DIRECTIONS:

Prior to application, examine orchard floor to locate trails and runways systems to be treated.

APPLICATION DIRECTIONS:

Hand Spot Baiting: Place about 1-1/2 oz. of bait in each active hole, trail, or runway at each tree site (10 lbs/acre). Cover each placement with grass or shingle to avoid exposing nontarget organisms.

Ground & Aerial Broadcasting: For aerial application only in those states specified above. For ground application use a commercially-made seed or fertilizer spreader. Uniformly apply 20 lb of bait per acre for Pine Voles and 10 lb of bait per acre for Meadow Voles. To avoid exposing nontarget organisms, clean up any bait in piles and return bait to application equipment or dispose of as specified under "Pesticide Disposal."

POST-APPLICATION INSTRUCTIONS:

If populations are high, make a second application 1-2 months after the first application.

NUT ORCHARDS (PECANS, ALMONDS, WALNUTS, FILBERTS/HAZELNUTS) FOR PINE AND MEADOW VOLE CONTROL

USE RESTRICTIONS: For control of pine and meadow voles in Nut Orchards in the following states:

Idaho Ohio Montana

Oregon Washington

Timing: Apply after fall harvest (including drops), before new spring growth, and when three days of good weather are expected. For other USE RESTRICTIONS, PRE-APPLICATION, APPLICATION, AND POST-APPLICATION DIRECTIONS, see use directions for pine and meadow voles in fruit orchards above.

CHRISTMAS TREE FARMS, COMMERCIAL NURSERIES, AND TREE PLANTATIONS FOR PINE AND MEADOW VOLE CONTROL

USE RESTRICTIONS: For control of pine and meadow voles in Christmas Tree Farms, Commercial Nurseries, and Tree Plantations in the following states:

Georgia Michigan Missouri North Carolina South Carolina Washington

Timing: Apply when three days of good weather are expected. For other USE RESTRICTIONS, PRE-APPLICATION, APPLICATION, AND POST-APPLICATION DIRECTIONS, see use directions for pine and meadow voles in fruit orchards above.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. *Household Label*

Pesticide Storage: Store in original container in a cool, dry place inaccessible to children and pets.

If empty: Nonrefillable container. Do not reuse this container. Place in trash or offer for recycling if available. **If partially filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any

Non-Household Label

indoor or outdoor drain.

Pesticide Storage: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Wastes resulting from the use of this product, and spoiled and unconsumed bait, may be disposed of on site or at an approved waste disposal facility.

Pesticide Container Disposal: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling is not available then dispose of empty container in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

Manufacturer and Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Manufacturer, and Buyer and User assume the risk of any such use. MANUFACTURER AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE **FOR** ANY INCIDENTAL, CONSEQUENTIAL OR **SPECIAL** DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF **MANUFACTURER** OR SELLER. THE REPLACEMENT OF THE PRODUCT.

Manufacturer and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of Manufacturer.

For non-emergency (e.g., current product information) call: 1-800-621-8829

Manufactured By: HACCO, Inc. 110 Hopkins Drive Randolph, WI 53956 U.S.A.

Ramik® is a Registered Trademark of HACCO, Inc

Code (NE.NO.) RN.NO.

Item No.: 0033370 Net Contents: 50 lbs.

UPC: 023626 00100 (8)

Format: UPC-A

Label:

This specimen label is intended for use only as a guide in providing general information regarding the directions, warning and cautions associated with the use of this product. As with any pesticide, always follow the label instructions on the package before using.

Ramik Brown/Page 3 of 3