

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

Lime Sulfur ULTRA

EPA Reg. No. 71096-11

For the suppression of *Pseudomonas Syringae* in Cherries

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. Follow all applicable directions, restrictions, precautions and Conditions of Sale on the EPA registered label. Refer to the container label for additional instructions. Always read and follow label directions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CHERRIES (SWEET AND TART) for the suppression of *Pseudomonas syringae* oozing/gum known as bacterial canker or bacterial blast in Cherries.

Fall Clean-up

Not for post-harvest application to harvested fruit. Apply 2 to 3 gallons of product per 100 gallons of water at 100 to 300 gallons of dilute spray per acre anytime after leaf drop begins. **DO NOT** make more than 2 applications total.

Dormant

Apply 2 to 3 gallons of product per 100 gallons of water at 100 to 300 gallons of dilute spray per acre. **DO NOT** make more than 2 applications.

Delayed Dormant

Apply 2 to 3 gallons of product per 100 gallons of water at 100 to 300 gallons of dilute spray per acre. **DO NOT** make more than 2 applications.

USE RESTRICTIONS: DO NOT apply more than 9 gallons (25.7 lbs. of Calcium Polysulfide) per application per acre or more than 75 gallons of LIME SULFUR ULTRA (214.5 lbs. of Calcium Polysulfide) per acre per year. **DO NOT** make more than 24 applications per year when applied at reduced use rates. Minimum Retreatment Interval = 7 days. Preharvest interval = 0 days.

Manufactured by:
OR-CAL, INC.
29454 Meadowview Rd.
Junction City, OR 97448

Dated 2/12/2024