



ALMOND ORCHARD SPRAYS

LIME SULFUR ULTRA

In Season Application

**Alternaria Leaf Spot, Powdery Mildew,
Rust, Scab, Shot Hole;**

Apply $\frac{1}{2}$ to $\frac{3}{4}$ gallon of product per 100 gallons of water. No more than 4 applications during growing season.

For Rust and Alternaria leaf spot, applications need to be made at the first sign of disease, before leaf symptoms appear, in late spring through summer at 10 to 14 day intervals. Non-Bearing Almonds may receive applications at 7 day intervals.

Less Bridging

**Little Known Resistance to
Pests/Diseases**

OMRI Listed

**More Effective than
Traditional Lime Sulfur**

Formulated with Potassium

**No Known Deleterious
Effects on Honey Bees**

**Post Reaction Leaves
Calcium Sulfate**



Thomas Putzel
Central & Southern CA, AZ
559-285-2122
Thomas@ORCALinc.com

Ruz Shokrollahi
Southeast US, TX
615-960-8188
Ruz@ORCALinc.com

Roger Haffner
OR, WA, ID, UT, MI, South FL
541-974-1712
Roger@ORCALinc.com

Jared Hajik
Northern CA
559-393-0691
Jared@ORCALinc.com

Scott Hansel
Northern CA
530-713-2911
ScottH@ORCALinc.com