

VINEYARD SPRAYS

LIME SULFUR ULTRA

In Season Application

From bud break through growing season application

Anthracnose

Apply ½ to 1 gallon of product per 100 gallons of water at 100 to 300 gallons of dilute spray per acre. Apply when new shoots are 4 to 6 inches long.

Phomopsis Cane and Leaf Spot

Apply ½ to 1 gallon of product per 100 gallons of water at 100 to 300 gallons of dilute spray per acre.

Powdery Mildew (Both A and B isolate)

Apply ½ to 1 gallon of product per 100 gallons of water. Apply a minimum of 100 gallons of dilute spray per acre. Repeat once or twice at 2 week intervals or when new shoots are a minimum of 4 to 6 inches long. Dilute sprays of 3 gallons of product per 100 gallons of water per acre may be applied when buds are swelling, when leaves are half grown and just before blossoming. If bronzing occurs on table grapes, increase spray dilution to ½ gallon per 125 gallons of water.



**LIME SULFUR
ULTRA**

*ALWAYS FOLLOW LABEL INSTRUCTIONS

Less Bridging

**Little Known Resistance
to Pests/Diseases**

OMRI Listed

**More Effective than
Traditional Lime Sulfur**

3% Formulated with Potassium

**No Known Deleterious
Effects on Honey Bees**

**Post Reaction Leaves
Calcium Sulfate**