



VINEYARD SPRAYS

LIME SULFUR ULTRA

In Season Application

From bud break through growing season application

Anthraco*nose*

Apply ½ to 1 gallon of product per 100 gallons of water at 100 to 300 gallons of dilute spray per acre. Apply when new shoots are 4 to 6 inches long.

Phomopsis Cane and Leaf Spot

Apply ½ to 1 gallon of product per 100 gallons of water at 100 to 300 gallons of dilute spray per acre.

Powdery Mildew (Both A and B isolate)

Apply ½ to 1 gallon of product per 100 gallons of water. Apply a minimum of 100 gallons of dilute spray per acre. Repeat once or twice at 2 week intervals or when new shoots are a minimum of 4 to 6 inches long. Dilute sprays of 3 gallons of product per 100 gallons of water per acre may be applied when buds are swelling, when leaves are half grown and just before blossoming. If bronzing occurs on table grapes, increase spray dilution to ½ gallon per 125 gallons of water.

 **LIME SULFUR
ULTRA**

*ALWAYS FOLLOW LABEL INSTRUCTIONS

Less Bridging

Little Known Resistance
to Pests/Diseases

OMRI Listed

More Effective than
Traditional Lime Sulfur

3% Formulated with Potassium

No Known Deleterious
Effects on Honey Bees

Post Reaction Leaves
Calcium Sulfate



For more information contact your Crop Advisor or visit ORCALinc.com

 **541-689-4413**
 29454 Meadowview Rd.
Junction City, OR 97448

**Scan to Find a
Rep in Your Area**

